ABDOMEN	
Location	□ Left □ Right □ Both □ Center
Pain Ratings	□ 0 □1 □2 □3 □4 □5 □6 □7 □8 □9 □10 (Excruciating)
Frequency	☐ Infrequent < 25% ☐ Occasional 25% to 50%
	☐ Frequent 50% to 75% ☐ Constant > 75%
Pain/Severity	□ No Pain □ Pain □ Numbness □ Tingling / □ Mild □ Moderate □ Severe
Associated with	☐ Bloating ☐ Constipation ☐ Diarrhea ☐ Nausea ☐ Rigidity
	□ Paresthesia □ Bruises □ Skin changes □ Vomiting
Radiates to	☐ Right Chest ☐ Left Chest ☐ Right Lower back ☐ Left Lower back ☐ Right Groin
	□ Left Groin □ Right Thigh □ Left Thigh
Described as	☐ Aching ☐ Dull ☐ Sharp ☐ Stabbing ☐ Throbbing
At its worst	☐ Morning ☐ Afternoon ☐ Evening ☐ Night
	☐ After Light Activities ☐ After Moderate Activities
What makes it better?	□ Medication □ Lying Down □ Standing □ Sitting
	□ Stretching □ Range of Motion □ Nothing
What makes it worse?	□ Palpation □ Bending □ Lifting □ Walking