BUTTOCKS	
Location	□ Right Buttock □ Left Buttock □ Both Buttocks
Pain Ratings	□ 0 □1 □2 □3 □4 □5 □6 □7 □8 □9 □10 (Excruciating)
Frequency	☐ Infrequent < 25% ☐ Occasional 25% to 50%
	☐ Frequent 50% to 75% ☐ Constant > 75%
Pain/Severity	□ No Pain □ Pain □ Numbness □ Tingling / □ Mild □ Moderate □ Severe
Associated with	□ Decreased ROM □ Increased Sensitivity □ Loss of Balance □ Soreness
Radiates to	☐ Left Lower back ☐ Right Lower back ☐ Right Mid-back ☐ Left Mid-back
	☐ Right Knee ☐ Left Knee ☐ Right Calf ☐ Left Calf
	□ Right Thigh □ Left Thigh
Described as	□ Aching □ Dull □ Sharp □ Stabbing □ Throbbing
At its worst	☐ Morning ☐ Afternoon ☐ Evening ☐ Night
	☐ After Light Activities ☐ After Moderate Activities
What makes it better?	☐ Medication ☐ Lying Down ☐ Standing ☐ Sitting
	□ Stretching □ Range of Motion □ Nothing
What makes it worse?	☐ Bending ☐ Extension ☐ Flexion ☐ Lying Down ☐ Walking Downstairs
	□ Walking Upstairs