| JAW (TMJ) | | | | | |
|-----------------------|--|------------------|--------------|-----------------|--------------|
| Location | □ Left | □ Right | □ Both | | |
| Pain Ratings | □0 □1 □2 | □3 □4 □5 | □6 □7 □8 | □9 □10 (Excruci | ating) |
| Frequency | ☐ Infrequent < 25% ☐ Occasional 25% to 50% | | | | |
| | □ Fred | quent 50% to 75% | % □ Coi | nstant > 75% | |
| Pain | □ No Pain | □ Pain | □ Numbness | □ Tingling | |
| Severity | □ Mild | □ Moderate | □ Severe | | |
| Associated with | □ Clicking | □ Popping | ☐ Grinding | □ Locking | □ Tightness |
| Radiates to | □ Head | □ Neck | ■ Ears | □ Face | |
| Described as | ☐ Aching | □ Dull | ☐ Sharp | □ Stabbing | ☐ Throbbing |
| At its worst | ☐ Morning | ☐ Afternoon | □ Evening | □ Night | |
| | After Activities | : □ Light | ■ Moderate | | |
| What makes it better? | □ Medication | ☐ Lying Dov | vn □ Standin | ng 🗆 Sitting | □ Stretching |
| | □ Range of Motion □ Nothing | | | | |
| What makes it worse? | □ Chewing | □ Yawning | □ Opening | mouth Clos | sing mouth |